

Quantum AetherDynamics Institute (QADI)

Filamentary Chronogeometry and Large-Scale Aether Organization: A Predictive Theory of Temporal-Torque Corridors and Cosmic Filament Structure

David W. Thomson III
Quantum AetherDynamics Institute (QADI)
Alma, Illinois, USA
ORCID: 0000-0002-5830-5427

Version 1.0 – June 20, 2026

This article is published by the Quantum AetherDynamics Institute (QADI).

It develops the Aether Physics Model (APM)

Abstract

The previous papers in the Temporal Torque Memory and Aether Organization series established that organized temporal-torque domains produce residual cosmic birefringence signatures that can be reconstructed numerically through Aether tomography and harmonic analysis. In this work, the framework is extended from reconstruction methodology to physical cosmology.

A temporal-torque density field,

$$\Omega_t(\mathbf{x}),$$

is introduced as a measure of organized five-dimensional chronogeometric structure. The gradients of this field generate preferred chronotorsional transport pathways that naturally organize into filamentary corridors. Cosmic filaments are shown to arise as integral curves of temporal-torque flow, while superclusters correspond to critical points of the temporal-torque landscape and cosmic voids emerge as minima of organized chronogeometric density.

A variational principle is developed in which filamentary structures extremize accumulated temporal torque. The resulting field equations predict anisotropic matter transport, preferred filament orientations, residual birefringence corridors, persistent handedness statistics, and characteristic filament spacing scales. Chronotorsional holonomy is shown to permit twisted filament bundles and large-scale helical organization.

The theory transforms temporal-torque memory from an observational signature into a predictive framework for the organization of the cosmic web and provides a set of falsifiable predictions that can be tested using future cosmic microwave background polarization surveys and large-scale structure observations.

Keywords: Temporal Torque; Chronogeometry; Cosmic Filaments; Cosmic Web; Aether Organization; Chronotorsional Holonomy; Cosmic Birefringence; Large-Scale Structure

1 Introduction

The previous papers of this series established a progressive framework for organized five-dimensional chronogeometric structure.

Paper I introduced temporal torque memory and demonstrated that organized Aether domains produce isotropic cosmic birefringence.

Paper II showed that residual birefringence maps may be interpreted as inverse tomographic probes of organized chronogeometric structure.

Paper III introduced quantitative definitions of temporal-torque density and chronotorsional holonomy.

Paper IV developed numerical methods for reconstructing organized domains using HEALPix G'orski et al. [2005] simulations and harmonic analysis Thomson III [2026d,a,c,b].

The present work extends the framework from numerical reconstruction to physical cosmology by investigating the organization of matter within temporally ordered Aether domains.

2 Temporal-Torque Density as an Order Parameter

The temporal-torque density field,

$$\Omega_t(\mathbf{x}),$$

is interpreted as an emergent macroscopic order parameter describing the degree of organized chronogeometric memory present within the Aether.

Regions possessing large values of

$$\Omega_t$$

exhibit highly coherent temporal organization, whereas regions of small

$$\Omega_t$$

correspond to weakly organized chronogeometric domains.

In this sense,

$$\Omega_t$$

plays a role analogous to magnetization in ferromagnetic systems or condensate density in superfluid systems. It is not introduced as an arbitrary scalar field but as a coarse-grained measure of persistent temporal organization.

3 Persistence of Temporal Memory

Temporal memory implies persistence.

Persistence implies spatial correlation.

Spatial correlation requires continuity of organized chronogeometric information.

Suppose organized temporal memory minimizes information discontinuities according to

$$E = \int (\nabla \Omega_t)^2 \, dV.$$

The Euler equation becomes

$$\nabla^2 \Omega_t = 0.$$

Solutions of Laplace's equation organize naturally into continuous streamlines connecting extrema of temporal organization.

These streamlines form corridor-like structures.

Cosmic filaments are therefore interpreted as the large-scale trajectories of persistent temporal memory within the organized Aether.

4 Chronogeometric Coherence Functional

The persistence principle may be written as a coherence functional for organized temporal memory,

$$E_\Omega = \int (\nabla \Omega_t)^2 \, dV.$$

This functional penalizes abrupt spatial discontinuities in temporal-torque density. It therefore represents the tendency of organized chronogeometric memory to maintain continuity across extended domains.

The corresponding Euler equation is

$$\nabla^2 \Omega_t = 0.$$

This result does not imply that the Universe is everywhere uniform. Rather, it implies that between sources, sinks, defects, and domain boundaries, organized temporal memory relaxes into continuous corridor-like structures.

The filamentary cosmic web is therefore interpreted as the large-scale solution structure of persistent chronogeometric coherence.

5 Temporal-Torque Flow

Define the temporal-torque flow vector by

$$\mathbf{F}(\mathbf{x}) = -\nabla \Omega_t(\mathbf{x}).$$

The flow vector points in the direction of maximum decrease of temporal-torque density.

The integral curves satisfy

$$\frac{d\mathbf{x}}{ds} = -\nabla \Omega_t.$$

These curves define preferred chronogeometric transport pathways.

Definition 5.1. A cosmic filament is an integral curve of the temporal-torque flow field.

6 Compatibility with Gravitational Structure Formation

The present framework does not replace gravitational structure formation.

Instead, organized temporal-torque domains establish preferred large-scale pathways of geometric coherence upon which gravitational evolution subsequently proceeds.

The sequence becomes

Temporal Organization → Coherent Corridors → Gravitational Amplification → Observed Cosmic Web.

Within this interpretation, gravitational instability determines how matter accumulates, while temporal-torque organization determines where coherent accumulation preferentially occurs.

Newtonian gravity, relativistic gravity, and gravitational torque theory therefore emerge as effective dynamical descriptions operating within a pre-organized chronogeometric framework.

7 Chronogeometric Pre-Structure and Gravitational Amplification

The present theory separates cosmic-web formation into two coupled stages.

The first stage is chronogeometric pre-structure. Organized temporal-torque memory establishes coherent pathways described by

$$\mathbf{F} = -\nabla\Omega_t.$$

The second stage is gravitational amplification. Matter responds to both the temporal-torque organization field and the gravitational potential,

$$\ddot{x}_i = -\partial_i\Omega_t - \partial_i\Phi_G.$$

Thus, gravity remains fully active. It determines accretion, collapse, orbital motion, and local dynamical evolution. The temporal-torque field determines the large-scale organization of preferred pathways through which gravitational dynamics become coherent.

This gives the theory a direct compatibility relation:

$$\text{Chronogeometric Pre-Structure} + \text{Gravitational Dynamics} = \text{Observed Cosmic Web}.$$

8 Variational Principle

The temporal-torque and gravitational fields are treated as an effective potential landscape through which matter propagates.

Consider the action

$$\mathcal{L} = \frac{1}{2}\dot{\mathbf{x}} \cdot \dot{\mathbf{x}} - \Omega_t(\mathbf{x}) - \Phi_G(\mathbf{x}).$$

$$S = \int \left[\frac{1}{2}\dot{x}_i\dot{x}_i - \Omega_t - \Phi_G\right] dt.$$

The Euler-Lagrange equations become

$$\frac{d}{dt}\left(\frac{\partial \mathcal{L}}{\partial \dot{x}_i}\right) - \frac{\partial \mathcal{L}}{\partial x_i} = 0,$$

giving

$$\ddot{x}_i = -\partial_i \Omega_t - \partial_i \Phi_G.$$

Matter therefore responds simultaneously to the organized temporal-torque landscape and the gravitational potential.

In regions where

$$|\partial_i \Omega_t| \gg |\partial_i \Phi_G|,$$

chronogeometric organization dominates the transport pathways.

In regions where

$$|\partial_i \Phi_G| \gg |\partial_i \Omega_t|,$$

gravitational dynamics dominate.

The observed cosmic web is interpreted as the coupled evolution of these two fields.

9 Coherent Gravitational Torques

Numerical studies of galaxy formation demonstrate that gravitational torques remain coherent over large spatial scales and guide matter flows into filamentary structures.

Within the present framework, this coherence is not accidental.

Persistent temporal memory establishes organized corridors that permit gravitational torques to add constructively over cosmological distances.

Gravitational torques therefore act as dynamical amplifiers of pre-existing chronogeometric organization.

10 Nodes, Filaments, Sheets, and Voids

Critical points satisfy

$$\nabla \Omega_t = 0.$$

Define the Hessian matrix Zel'dovich [1970], Forero-Romero et al. [2009], Sousbie [2011], Sousbie et al. [2011]

$$H_{ij} = \frac{\partial^2 \Omega_t}{\partial x_i \partial x_j}.$$

The eigenvalues of

$$H_{ij}$$

classify the morphology.

Three negative eigenvalues correspond to supercluster nodes.

Two negative eigenvalues correspond to filamentary structures.

One negative eigenvalue corresponds to sheet-like organization.

No negative eigenvalues correspond to void regions.

11 Chronotorsional Holonomy

Let

$$\tau_t$$

denote the chronotorsional holonomy.

Define

$$\mathbf{T} = \nabla \tau_t.$$

The generalized filament equation becomes

$$\mathbf{F} = -\nabla \Omega_t + \gamma \nabla \tau_t,$$

where

$$\gamma$$

is a coupling coefficient.

Regions satisfying

$$\nabla \times \mathbf{F} \neq 0$$

develop twisted filament bundles and intrinsic helicity.

The theory therefore predicts preferred filament handedness and large-scale helical organization.

12 Filament Spin and Handedness Conservation

Chronotorsional holonomy does not merely bend filamentary corridors. It also permits filament bundles to acquire persistent spin orientation and handedness.

Let a filament corridor be represented by an integral curve

$$\mathbf{x}(s),$$

with tangent vector

$$\hat{\mathbf{t}} = \frac{d\mathbf{x}/ds}{|d\mathbf{x}/ds|}.$$

The temporal-torque flow field is

$$\mathbf{F} = -\nabla\Omega_t + \gamma\nabla\tau_t.$$

The filament spin functional is defined by

$$\mathbf{S}_f = \oint \mathbf{r} \times \mathbf{F}\, ds,$$

where

$$\mathbf{r}$$

is the displacement vector measured from the local filament axis.

If

$$\mathbf{S}_f = 0,$$

the filament corridor possesses no persistent handedness. If

$$\mathbf{S}_f \neq 0,$$

the filament corridor possesses a preferred spin orientation.

The associated handedness scalar may be written as

$$h_f = \hat{\mathbf{t}} \cdot (\nabla \times \mathbf{F}) .$$

Using the generalized flow equation,

$$\nabla \times \mathbf{F} = -\nabla \times (\nabla \Omega_t) + \gamma \nabla \times (\nabla \tau_t) .$$

For smooth single-valued scalar fields,

$$\nabla \times \nabla \Omega_t = 0$$

and

$$\nabla \times \nabla \tau_t = 0.$$

However, chronotorsional holonomy is not required to be globally single-valued. In the presence of organized topological structure, domain boundaries, or chronotorsional defects,

$$\nabla \times \nabla \tau_t \neq 0.$$

Therefore,

$$h_f = \gamma \hat{\mathbf{t}} \cdot (\nabla \times \nabla \tau_t) .$$

This expression shows that filament handedness is generated by nontrivial chronotorsional holonomy.

The condition

$$\nabla \times \nabla \tau_t \neq 0$$

is interpreted as evidence of nontrivial topological organization.

Chronotorsional holonomy therefore behaves analogously to a multi-valued phase field possessing defects, branch surfaces, or noncontractible loops.

In such regions, the usual vector identity

$$\nabla \times \nabla \tau_t = 0$$

fails globally even though it remains valid locally away from topological defects.

Filament handedness therefore arises from organized chronotorsional topology rather than from local vector calculus alone.

The sign of

$$h_f$$

distinguishes right-handed and left-handed filament organization. The magnitude of

$$h_f$$

measures the strength of the filament's helical chronogeometric structure.

Persistent temporal memory implies that filament handedness should not be purely random. If the Aether retains organized chronotorsional memory across cosmological distances, then neighboring filaments should display correlated spin orientation, correlated galaxy angular momenta, and correlated residual birefringence signatures.

This leads to a distinctive prediction:

$$\mathrm{corr}\,(h_f, \delta\beta) \neq 0.$$

Filaments with nonzero handedness should therefore exhibit corresponding polarization-rotation structure in residual birefringence maps.

Within this framework, filament spin is not merely a consequence of local gravitational accretion. Gravitational torques amplify and populate the filament, but chronotorsional holonomy establishes the persistent handedness structure through which matter flows.

Thus, large-scale filament spin, galaxy-spin alignments, and residual birefringence handedness are interpreted as mutually related expressions of organized temporal-torque memory Peebles [1969], White [1984], Codis et al. [2018], Wang et al. [2021].

13 Preferred Filament Spacing

Assume that organization minimizes the functional

$$E = \int \left[-(\nabla\Omega_t)^2 + L^2(\nabla^2\Omega_t)^2 \right] dV.$$

Variation yields

$$\nabla^2\Omega_t + L^2\nabla^4\Omega_t = 0.$$

$$L = L(\Omega_t, \tau_t).$$

$$\lambda_c = 2\pi L(\Omega_t, \tau_t).$$

The theory predicts that filament spacing should correlate with the degree of temporal organization and chronotorsional memory.

Fourier solutions satisfy

$$k^2(1 - L^2k^2) = 0.$$

The characteristic scale becomes

$$k_c = \frac{1}{L},$$

giving

$$\lambda_c = 2\pi L.$$

The theory therefore predicts preferred filament spacing scales.

The simplest coherence functional,

$$E_\Omega = \int (\nabla\Omega_t)^2 dV.$$

produces a correlation length over which temporal organization remains spatially coherent.

When higher-order curvature effects become significant, the functional

$$E = \int \left[-(\nabla\Omega_t)^2 + L^2(\nabla^2\Omega_t)^2\right] dV$$

permits periodic organization with characteristic scale

$$\lambda_c = 2\pi L.$$

The theory therefore allows both continuously correlated filament corridors and preferred filament spacing scales.

14 Filaments as Fossilized Temporal Memory

If temporal organization persists through the Aether, its gradients necessarily produce coherent pathways that connect regions of differing chronogeometric density.

Matter subsequently accumulates along these pathways through gravitational amplification.

The observed cosmic web may therefore be interpreted as a fossil record of organized temporal memory.

Cosmic filaments are not merely gravitational accidents.

They are observable trajectories of persistent chronogeometric organization.

15 Residual Birefringence Corridors

Residual birefringence is assumed to depend upon organized temporal torque according to

$$\delta\beta = f(\Omega_t, \tau_t).$$

To first order,

$$\delta\beta \propto |\nabla\Omega_t| + \eta\tau_t.$$

The theory predicts:

1. filamentary birefringence corridors,
2. coherent polarization ridges,
3. preferred filament orientations,
4. handed polarization structures,
5. anisotropic multipole signatures.

16 Observational Predictions

The theory developed in this paper produces multiple observational predictions. These predictions are not mutually exclusive. They represent different observational projections of the same organized chronogeometric structure.

16.1 Filamentary Birefringence Corridors

Residual birefringence maps should exhibit elongated corridor-like features aligned with large-scale cosmic filaments.

To first order,

$$\delta\beta \propto |\nabla\Omega_t| + \eta\tau_t.$$

Therefore, regions of strong temporal-torque gradient and nontrivial chronotorsional holonomy should produce coherent polarization-rotation signatures.

16.2 Correlation Between Birefringence and Cosmic-Web Filaments

Observed cosmic-web filaments should correlate statistically with gradients in reconstructed residual birefringence maps.

The predicted correlation is

$$\mathrm{corr}\left(\hat{\mathbf{f}}, \nabla\delta\beta\right) \neq 0,$$

where

$$\hat{\mathbf{f}}$$

is the observed filament orientation.

16.3 Supercluster Nodes as Temporal-Torque Extrema

Superclusters should coincide with critical points of the temporal-torque density field.

These points satisfy

$$\nabla \Omega_t = 0$$

and

$$\det(H_{ij}) \neq 0.$$

The eigenvalue structure of

$$H_{ij}$$

determines whether the critical point behaves as a node, saddle, or transition structure.

16.4 Voids as Minima of Organized Temporal Memory

Cosmic voids should correspond to regions of weak organized temporal-torque density.

In void interiors,

$$\Omega_t \to \Omega_{\min},$$

and

$$|\nabla \Omega_t| \to 0.$$

The theory therefore predicts reduced filament density, weak residual birefringence, and minimal chronotorsional handedness within mature void regions.

16.5 Matter-Flow Anisotropy

Matter flows should preferentially align with the combined chronogeometric and gravitational acceleration field,

$$\ddot{x}_i = -\partial_i \Omega_t - \partial_i \Phi_G.$$

When gravitational amplification dominates the local matter distribution, the temporal-torque term remains observable through large-scale anisotropic alignment.

Thus, the theory predicts that matter transport should retain a statistical memory of

$$-\nabla \Omega_t.$$

16.6 Coherent Gravitational Torque Alignment

Gravitational torques should add constructively along temporal-torque corridors.

Therefore, galaxy angular momentum, filament spin, and matter-flow orientation should show enhanced coherence along filamentary corridors compared with randomly selected regions of equal matter density.

16.7 Persistent Filament Handedness

Filament handedness statistics should depart from isotropy.

The handedness scalar

$$h_f = \hat{\mathbf{t}} \cdot (\nabla \times \mathbf{F})$$

should exhibit persistent sign coherence over large spatial domains when chronotorsional holonomy is organized.

The prediction is

$$\langle h_f \rangle_{\text{domain}} \neq 0$$

for coherent chronotorsional domains.

16.8 Correlation Between Filament Handedness and Residual Birefringence

Because residual birefringence depends upon organized temporal torque and chronotorsional holonomy, filament handedness should correlate with polarization rotation.

The predicted relation is

$$\text{corr}\,(h_f, \delta\beta) \neq 0.$$

This is a particularly important prediction because it links large-scale matter organization to polarization topology.

16.9 Preferred Filament Spacing

The characteristic filament spacing should be related to the correlation length of organized temporal memory,

$$\lambda_c = 2\pi L(\Omega_t, \tau_t).$$

Therefore, filament spacing should not be purely accidental. It should vary with the strength of temporal-torque organization and chronotorsional memory.

16.10 Filament Connectivity and Holonomy

The number of filaments entering a supercluster node should correlate with the local holonomy structure.

Regions of strong chronotorsional organization should exhibit higher filament connectivity and more coherent angular structure around nodes.

16.11 Residual Birefringence as a Cosmic-Web Tracer

Residual birefringence should reconstruct portions of the cosmic web even where luminous matter is sparse.

This prediction follows because

$$\delta\beta$$

traces organized temporal-torque structure rather than luminous matter alone.

Therefore, residual birefringence maps may reveal hidden or weakly luminous filament corridors.

16.12 Multipole Alignment

Large-angle polarization multipoles should show nonrandom alignment with reconstructed temporal-torque corridors.

The predicted observational chain is

$$\Omega_t(\mathbf{x}) \to \text{filaments} \to \delta\beta(\theta, \phi) \to a_{\ell m} \to \text{observations}.$$

Thus, anisotropic multipole signatures should not be treated as isolated anomalies. They may represent harmonic projections of organized filamentary chronogeometry.

16.13 Persistence Across Cosmological Epochs

If temporal-torque memory is persistent, then filamentary organization should retain coherent structural signatures across cosmological time.

The theory predicts that high-redshift filament structures, galaxy-spin alignments, and residual birefringence templates should show continuity with lower-redshift cosmic-web organization.

16.14 Summary of Predictions

The principal predictions are:

1. residual birefringence maps should contain filamentary corridors;
2. birefringence gradients should correlate with observed cosmic-web filaments;
3. superclusters should correspond to extrema or saddle points of Ω_t;
4. cosmic voids should correspond to minima of temporal-torque density;
5. matter flows should retain alignment with $-\nabla\Omega_t$;
6. gravitational torques should add coherently along temporal-torque corridors;
7. filament handedness statistics should depart from isotropy;
8. filament handedness should correlate with residual birefringence;
9. preferred filament spacing should depend upon $L(\Omega_t, \tau_t)$;
10. filament connectivity should correlate with chronotorsional holonomy;
11. residual birefringence should trace cosmic-web structure even where luminous matter is sparse;
12. large-angle polarization multipoles should align with reconstructed temporal-torque corridors;
13. filamentary organization should persist across cosmological epochs.

Together, these predictions distinguish filamentary chronogeometry from a purely gravitational description while remaining fully compatible with gravitational amplification, orbital dynamics, and relativistic gravitational structure formation.

17 Distinguishing Observational Signatures

Several predictions of this framework overlap with gravitational cosmic-web morphology. This is expected because gravity amplifies the same large-scale structures that temporal-torque organization pre-structures.

The distinctive signatures arise where matter geometry and polarization topology are correlated.

The strongest distinguishing prediction is

$$\mathrm{corr}\,(h_f, \delta\beta) \neq 0.$$

A second distinguishing prediction is

$$\mathrm{corr}\left(\hat{\mathbf{f}}, \nabla\delta\beta\right) \neq 0.$$

A third distinguishing prediction is that residual birefringence should trace filamentary corridors even where luminous matter is weak or sparsely sampled.

Gravity predicts matter clustering.

Filamentary chronogeometry predicts matter clustering together with polarization rotation, handedness coherence, and residual birefringence corridors.

The observational distinction is therefore not whether filaments exist, but whether filament geometry, filament handedness, and cosmic polarization rotation share a common organized chronogeometric source.

Within the present framework, the cosmic web possesses three simultaneous observables:

$$\text{Matter Distribution}, \qquad \text{Filament Handedness}, \qquad \text{Polarization Topology}.$$

The theory predicts that these observables are statistically coupled because they originate from the same organized chronogeometric structure.

The simultaneous correlation of all three observables would constitute strong evidence for persistent temporal-torque organization within the Aether.

18 Parameterization and Future Quantitative Tests

The present paper develops the physical theory of filamentary chronogeometry at the effective-field level. The fields

$$\Omega_t(\mathbf{x})$$

and

$$\tau_t(\mathbf{x})$$

are treated as macroscopic order parameters describing organized temporal-torque density and chronotorsional holonomy.

A future quantitative implementation should determine these fields from reconstructed birefringence maps, galaxy filament catalogs, and large-scale structure surveys Minami and Komatsu [2020], Diego-Palazuelos et al. [2022], LiteBIRD Collaboration [2023], Abazajian et al. [2016].

The principal empirical quantities to estimate are

$$L(\Omega_t, \tau_t), \qquad \gamma, \qquad \eta,$$

where (L) controls filament spacing, (γ) controls holonomy-driven handedness, and (η) controls the birefringence response to chronotorsional holonomy.

The predicted observational chain is

$$\Omega_t(\mathbf{x}) \to \mathbf{F}(\mathbf{x}) \to \text{filament corridors} \to h_f \to \delta\beta \to a_{\ell m}.$$

This chain provides a direct path from theory to mock observations.

The next stage of the program is therefore to construct simulated catalogs in which temporal-torque corridors are generated from specified

$$\Omega_t$$

and

$$\tau_t$$

fields, projected into residual birefringence maps, and compared against observed cosmic-web filament catalogs.

This program will allow the theory to move from qualitative correlations to quantitative forecasts for filament spacing, handedness statistics, birefringence amplitude, and multipole alignment.

19 Discussion

Within this framework, the cosmic web is interpreted as the residual imprint of organized five-dimensional chronogeometric structure.

Matter does not merely collapse gravitationally within an otherwise featureless background. Instead, matter propagates along preferred chronogeometric pathways established by organized temporal-torque domains.

The resulting cosmic network represents an observable fossil record of large-scale Aether organization.

20 Conclusions

A predictive theory of filamentary chronogeometry has been developed in which organized temporal-torque density naturally generates filamentary corridors, sheets, voids, and supercluster nodes.

Chronotorsional holonomy produces twisted filament bundles and persistent handedness statistics.

Residual cosmic birefringence maps are predicted to contain filamentary signatures that directly encode organized five-dimensional chronogeometric structure.

The progression of the Temporal Torque Memory and Aether Organization series therefore becomes

$$\text{Memory} \to \text{Tomography} \to \text{Quantification} \to \text{Reconstruction} \to \text{Cosmic Organization}.$$

The cosmic web may therefore be interpreted as an observable manifestation of organized temporal-torque structure within the Aether.

Acknowledgments

The Quantum AetherDynamics Institute (QADI) is a private 501(c)(3) research institute developing the Aether Physics Model (APM) and Quantum Measurement Units (QMU) Thomson III [2025]. Publications and data are archived with DOIs in the AetherPhysics Zenodo community: zenodo.org/communities/aetherphysics.

References

K. N. Abazajian et al. CMB-S4 science book, first edition, 2016.

S. Codis, A. Jindal, N. E. Chisari, D. Vibert, Y. Dubois, C. Pichon, and J. Devriendt. Galaxy orientation with the cosmic web across cosmic time. *Monthly Notices of the Royal Astronomical Society*, 481(4):4753–4774, 2018. doi: 10.1093/mnras/sty2567.

P. Diego-Palazuelos, J. R. Eskilt, Y. Minami, M. Tristram, R. M. Sullivan, A. J. Banday, R. B. Barreiro, H. K. Eriksen, K. M. G'orski, R. Keskitalo, E. Komatsu, E. Mart'inez-Gonz'alez, D. Scott, P. Vielva, and I. K. Wehus. Cosmic birefringence from planck public release 4. *Physical Review Letters*, 128:091302, 2022. doi: 10.1103/PhysRevLett.128.091302.

J. E. Forero-Romero, Y. Hoffman, S. Gottl"ober, A. Klypin, and G. Yepes. A dynamical classification of the cosmic web. *Monthly Notices of the Royal Astronomical Society*, 396(3):1815–1824, 2009. doi: 10.1111/j.1365-2966.2009.14885.x.

K. M. G'orski, E. Hivon, A. J. Banday, B. D. Wandelt, F. K. Hansen, M. Reinecke, and M. Bartelmann. HEALPix: A framework for high-resolution discretization and fast analysis of data distributed on the sphere. *The Astrophysical Journal*, 622(2):759–771, 2005. doi: 10.1086/427976.

LiteBIRD Collaboration. Probing cosmic inflation with the litebird cosmic microwave background polarization survey. *Progress of Theoretical and Experimental Physics*, 2023(4):042F01, 2023. doi: 10.1093/ptep/ptac150.

Y. Minami and E. Komatsu. New extraction of the cosmic birefringence from the planck 2018 polarization data. *Physical Review Letters*, 125:221301, 2020. doi: 10.1103/PhysRevLett.125.221301.

P. J. E. Peebles. Origin of the angular momentum of galaxies. *The Astrophysical Journal*, 155:393, 1969. doi: 10.1086/149876.

T. Sousbie. The persistent cosmic web and its filamentary structure – i. theory and implementation. *Monthly Notices of the Royal Astronomical Society*, 414(1):350–383, 2011. doi: 10.1111/j.1365-2966.2011.18394.x.

T. Sousbie, C. Pichon, and H. Kawahara. The persistent cosmic web and its filamentary structure – ii. illustrations. *Monthly Notices of the Royal Astronomical Society*, 414(1):384–403, 2011. doi: 10.1111/j.1365-2966.2011.18395.x.

David W. Thomson III. Quantum aetherdynamics institute (qadi): Community doi overview and citation guide. Zenodo, 2025. URL https://doi.org/10.5281/zenodo.17479314.

David W. Thomson III. Aether tomography: Residual cosmic birefringence as a probe of organized five-dimensional chronogeometric structure, June 2026a. URL https://doi.org/10.5281/zenodo.20754696.

David W. Thomson III. Numerical reconstruction of organized temporal-torque domains and simulated residual cosmic birefringence maps, June 2026b. URL https://doi.org/10.5281/zenodo.20779504.

David W. Thomson III. Quantitative definition of temporal torque density and chronotorsional holonomy: A qmu derivation of organized five-dimensional chronogeometric structure, June 2026c. URL https://doi.org/10.5281/zenodo.20778298.

David W. Thomson III. Temporal torque memory and galactic-scale aether organization: A predictive framework for cosmic birefringence, June 2026d. URL https://doi.org/10.5281/zenodo.20753446.

P. Wang, N. I. Libeskind, E. Tempel, X. Kang, and Q. Guo. Possible observational evidence for cosmic filament spin. *Nature Astronomy*, 5:839–845, 2021. doi: 10.1038/s41550-021-01380-6.

S. D. M. White. Angular momentum growth in protogalaxies. *The Astrophysical Journal*, 286:38–41, 1984. doi: 10.1086/162573.

Ya. B. Zel'dovich. Gravitational instability: An approximate theory for large density perturbations. *Astronomy and Astrophysics*, 5:84–89, 1970.